

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/74	A1	(11) International Publication Number: WO 96/22780
		(43) International Publication Date: 1 August 1996 (01.08.96)
(21) International Application Number: PCT/US96/00633		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 24 January 1996 (24.01.96)		
(30) Priority Data:		
08/379,695 27 January 1995 (27.01.95) US		
08/440,417 12 May 1995 (12.05.95) US		
(71) Applicant: THE BOARD OF REGENTS OF THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 W. 7th Street, Austin, TX 78701 (US).		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(72) Inventors: LICHTENBERGER, Lenard, M.; 5427 Darnell Street, Houston, TX 77096 (US). BUTLER, Bruce, D.; 1609 Milford, Houston, TX 77006 (US).		
(74) Agent: MAYFIELD, Denise, L.; Akin, Gump, Strauss, Hauer & Feld, L.L.P., Suite 1900, 816 Congress Avenue, Austin, TX 78701 (US).		

(54) Title: METHODS OF ENHANCING THE THERAPEUTIC ACTIVITY OF NSAIDS AND COMPOSITIONS OF ZWITTERIONIC PHOSPHOLIPIDS USEFUL THEREIN



(57) Abstract

Disclosed are compositions comprising non-steroid anti-inflammatory drugs (NSAID's) complexed with zwitterionic, neutral phospholipids, or both, having reduced gastrointestinal irritating effects and enhanced anti-pyretic, analgesic, and anti-inflammatory activity. Also disclosed are improved methods of using the complexes for treating fever, inflammation, and preventing platelet aggregation. In some embodiments, the anti-pyretic activity of sub-therapeutically used amounts of NSAID's are enhanced to elicit anti-pyretic activity *in vivo* when associated (noncovalently) with zwitterionic phospholipids, such as dipalmitoyl phosphatidyl choline.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**METHODS OF ENHANCING THE THERAPEUTIC
ACTIVITY OF NSAIDS AND COMPOSITIONS OF
ZWITTERIONIC PHOSPHOLIPIDS USEFUL THEREIN**

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 The present invention relates generally to the fields of pharmacology and medicine and more particularly, it concerns the treatment and prevention of fever, pain and inflammation with non-steroidal anti-inflammatory drugs (NSAIDs) complexed with phospholipids, and in other embodiments in further combination with neutral lipids. The invention also provides methods for retarding platelet aggregation, and the application of these methods in treating cardiovascular and vascular diseases as it relates to platelet activity.

15

2. Description of the Related Art

20 The consumption of NSAIDs among the general populace is unparalleled by any other drug class due to their great efficacy in the treatment of pain, inflammation and fever (Rainsford, 1985). The widespread usage of these drugs is anticipated to increase even further due to their efficacy in the treatment of osteoarthritic and generalized aches and pain as the elderly increase as a percentage of the population (Alexander *et al.*, 1985; Jolobe and Montgomery, 1984), and as NSAIDs are employed in the treatment/prevention of stroke and cardiovascular disease.

25

The major concern with these developments relates to the tendency of NSAIDs to induce gastrointestinal (GI) mucosal lesions, perforations and bleeding resulting in significant morbidity and mortality, even in occasional NSAID users (Rainsford, 1989; Graham, 1989; Allison *et al.*, 1992). Strategies to reduce the gastroduodenal injurious effects of these drugs with enteric coatings, have had limited success due to the delayed therapeutic actions of these specially packaged NSAIDs (Alpsten *et al.*, 1982; Mojaverian *et al.*, 1987).

30

SUBSTITUTE SHEET (RULE 26)

Although it is clear that the GI side-effects of NSAIDs are in part attributable to their ability to inhibit the biosynthesis of gastroprotective prostaglandins, a significant amount of evidence exists that NSAIDs act locally on the mucosa to induce GI ulcers and bleeding by a prostaglandin-independent mechanism (Rainsford, 1989; Whittle *et al.*, 1980; Whittle, 1981; Ligumsky *et al.*, 1982; McCormack and Brune, 1987).

The present inventor and others have obtained evidence that the mucosa of the stomach and other regions of the GI tract have hydrophobic, non-wettable properties, that protect the underlying epithelium from gastric acid and other luminal toxins (Hills *et al.*, 1983; Goddard *et al.*, 1987; Goddard *et al.*, 1990; Kao *et al.*, 1990). This biophysical characteristic, which can be quantified by contact angle analysis, appears to be attributable to the presence of an extracellular lining of surfactant-like phospholipid on the luminal aspects of the mucus gel layer (Goddard *et al.*, 1990; Kao *et al.*, 1990). Evidence has also come forth that these zwitterionic phospholipids are synthesized in surface mucus cells of the stomach, as well as those present in discrete submucosal glands of the GI tract, where they are stored in specific organelles and secreted by a prostaglandin-dependent pathway (Kao and Lichtenberger, 1991). It has also been reported that aspirin and other NSAIDs have the ability to rapidly transform the gastric mucosa from a non-wettable to a wettable state within minutes after luminal administration, thereby increasing the tissue's susceptibility to the corrosive actions of gastric acid (Hills *et al.*, 1983; Goddard *et al.*, 1987; Goddard *et al.*, 1990; Kao *et al.*, 1990).

One solution to this problem has been to formulate injectable solutions of NSAIDs and thus bypass the GI tract completely. The low water solubility of these drugs, however, has caused problems with this technique. Stable, injectable solutions of indoleacetic and indanacetic acid derivatives have been developed to address this problem, by complexing these NSAIDs with phosphatidylcholine and phosphatidylethanolamine derivatives (See US Patent 4,309,420).

US Patent 4,421,747 describes NSAIDs complexed with phospholipids for oral administration. These complexes were shown to retain their anti-inflammatory action and to have reduced ulcer formation in rats. However, no enhancement of therapeutic effects was reported with these preparations.

5

WO 91/16920 (Vical Inc.) relates to phospholipid prodrug derivatives of a salicylate or non-steroidal, anti-inflammatory drug. These preparations are made by combining salicylic acid or NSAID with a phospholipid in the presence of a coupling agent, thereby producing a covalently linked NSAID-phospholipid compound. These prodrugs are described as useful in reducing the toxicity of high dose, long term usage of NSAID preparations.

10

JP 3176425 (Nippon Shinyaku KK) relates to compositions including non-steroidal, anti-inflammatory drugs together with neutral lipids and phospholipids in a fat and oil emulsion. Although the method of preparation is not described in the abstract, these compositions appear to be encapsulated in lipid, such as in a micelle. The combination of the drug with the neutral lipids and the phospholipids is described as not affecting the drug's pharmacological actions.

15

JP 63048228 (Toa Eiyo KK) relates to topically applied compositions that include non-steroidal anti-inflammatory drug together with phospholipid and a "disintegrator". The disintegrator is described as providing for a preparation with improved dispersability and increased absorptivity. JP 63048226 (Ono Pharmaceutical KK) relates to compositions that include a phospholipid base (such as phosphatidylcholine) and an anti-inflammatory agent (such as acetylsalicylic acid and indomethacin). KK JP 58150508 (Ono Pharmaceutical) relates to topical compositions that include a phospholipid base (such as phosphatidylcholine) and an anti-inflammatory agent (such as acetylsalicylic acid and indomethacin).

20

25

30

US Patent 4,369,182 (Nattermann & CIE), relates to inflammation-preventing pharmaceutical compositions for oral administration. The compositions are prepared

and then lyophilized into powder form. The described compositions include natural and synthetic phospholipids (dipalmitoylphosphatidylcholine (DPPC)), in combination with nonsteroidal agents including salicylic acid, acetyl-salicylic acid, diflunisal, indomethacin, glucametacine, acemetacin, sulindac, ibuprofen, naproxen, tolmetin and other NSAID's. Also described are NSAID's in combination with phosphatidylcholine preparations, named phospholipons. US Patent 4,421,747 relates to methods of alleviating inflammation with compositions as described in the '182 patent.

Despite the extensive work in the area of NSAIDs, a need continues to exist in the art for preparations that include reduced amounts of this useful class of drug without loss of therapeutic efficacy. Methods and compositions that provide for similar or enhanced anti-pyretic, anti-inflammatory, anti-platelet and analgesic activity at lower doses than currently prescribed for pharmacological activity would also render this very valuable class of drugs available to those previously unable to tolerate standard and/or prolonged therapeutic regimens of NSAID.

SUMMARY OF THE INVENTION

The present invention seeks to overcome these and other drawbacks inherent in the prior art by providing compositions and methods capable of maintaining and/or improving the pharmacological activity of non-steroidal anti-inflammatory drugs by noncovalent association with zwitterionic phospholipids. In some embodiments, these preparations may further include neutral lipids, such as the triglycerides.

The present invention illustrates the ability of one or more zwitterionic phospholipids to enhance the fever-reducing potential of an NSAID. Pharmacological activity of low dose NSAID to reduce inflammation and pain may also be observed, and in some cases enhanced by chemically associating the NSAID with zwitterionic phospholipid, such as phosphatidyl choline (PC), dipalmitoylphosphatidylcholine (DPPC), and other disaturated phosphatidyl cholines, and the like. In some embodiments, the association of NSAID and zwitterionic phospholipid is of a non-

covalent nature. In still other embodiments, the NSAID and zwitterionic phospholipid compositions may be further described as including more or less equimolar amounts of these ingredients. The compositions may also, of course, comprise a pharmaceutically acceptable carrier in any form, such as solid powder, gel or liquid form.

The present invention focusses techniques that are demonstrated in some cases to enhance the therapeutic activities of NSAIDs. This is accomplished without the disadvantage of hindering the pharmacological activity or therapeutic bioavailability of the drug rendering the preparations effective even at low doses.

Accordingly, the present invention provides in one aspect a method for enhancing the antipyretic activity of a nonsteroidal anti-inflammatory drug (NSAID). This method comprises providing a non-covalently associated combination of a zwitterionic phospholipid with an amount of a nonsteroidal anti-inflammatory drug that provides reduced anti-pyretic activity in the absence of the zwitterionic phospholipid. Some embodiments of the present preparations are further described as being essentially free of anionic phospholipid, or as including an amount of anionic phospholipid that is biologically inert and/or not an active component, of the preparation.

The term "essentially free" as used in the description of the present invention, is understood to mean compositions that include less than about 0.10% of anionic phospholipid, and in even further defined embodiments, less than 0.01% anionic phospholipid. As used in the description of the present invention, the term zwitterionic phospholipid embraces a wide range of phospholipids, including but not limited to phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingomyelin and other ceramides, as well as various other zwitterionic phospholipids. In some embodiments, these compositions are essentially free of anionic phospholipid.

In other embodiments of the described method, the amount of nonsteroidal anti-inflammatory drug is defined as an amount that provides reduced antipyretic activity in the absence of the zwitterionic phospholipid. Such amounts of the drug sub-therapeutically effective amounts thereof. This activity or lack of activity is observed in the absence of zwitterionic phospholipid, while the same or about the same amount of the NSAID does demonstrate pharmacological activity in the presence of zwitterionic phospholipid. In this regard, the phenomenon is observed that the combination of low amounts of nonsteroidal anti-inflammatory drugs with phospholipid have potent pharmacological activity, while doses of the drug alone (i.e., without zwitterionic phospholipid) do not.

The present method employs compositions that may include any variety of those drugs generally classified as nonsteroidal anti-inflammatory drugs. By way of example, these drugs include ibuprofen, piroxicam, salicylate, aspirin, naproxen, indomethacin, diclofenac, or any mixture thereof. In particular embodiments, the nonsteroidal anti-inflammatory drug is salicylate. By way of example and not limitation, NSAID's useful in the practice of the invention, include those noted in Table 1.

TABLE 1

**Nonsteroidal Anti-Inflammatory Drugs To Be Used
in Combination with Zwitterionic Phospholipids**

5

10

15

20

25

30

Propionic acids	Fenoprofen calcium	Nalfon
	Flurbiprofen	Ansaid
	Suprofen	
	Benoxaprofen	
	Ibuprofen (prescription)	Motrin
	Ibuprofen (200 mg. over the counter)	Nuprin, Motrin 1B
	Ketoprofen	Orduis, Oruvall
	Naproxen	Naprosyn
	Naproxen sodium	Aleve, Anaprox, Aflaxen
	Oxaprozin	Daypro
Acetic acids	Diclofenac sodium	Voltaren
	Diclofenac potassium	Cataflam
	Etodolac	Lodine
	Indomethacin	Indocin
	Ketorolac tromethamine (intramuscular)	Acular, Toradol
	Ketorolac (oral)	Toradol
Ketones	Nabumetone	Relafen
	Sulindac	Clinoril
	Tolmetin sodium	Tolectin
Fenamates	Meclofenamate sodium	Meclofen
	Mefenamic acid	Ponstel
Oxicams	Piroxicam	Dolibid
Salicylic acid	Diflunisal	Feldene
	Aspirin	
Pyrazolin acid	Oxyphenbutazone	Tandearil
	Phenylbutazone	Butazolidin

SUBSTITUTE SHEET (RULE 26)

NSAIDs such as benoxaprofen, ketoprofen, oxaprozin, etodolac, ketorolac
tromethamine, ketorolac and nabumetone, together with zwitterionic phospholipid,
comprise still other particular embodiments of the invention, again both with and
5 without neutral lipid.

In particular embodiments, the zwitterionic phospholipid in the compositions
is dipalmitoyl phosphatidylcholine, phosphatidyl choline, or a mixture thereof.

10 The pharmacological, particularly anti-pyretic, activity of NSAIDs, is shown
to be enhanced several fold over preparations with similar doses without zwitterionic
phospholipid. Surprisingly, the present inventors have found that combination of the
aforedescribed nonsteroidal anti-inflammatory drugs with zwitterionic phospholipid
dramatically enhances the anti-pyretic activity and potency of the drug, even at sub-
15 therapeutically active doses, and in some cases from about 2-fold to about 6-fold
relative to non-phospholipid containing preparations. Hence, the methods are expected
to be particularly efficacious in reducing fever in a mammal having reduced tolerance
for NSAID's.

20 Amounts ranging between one-tenth and one-half that typically necessary to
illicit a fever-reducing response in a mammal may thus be realized employing the
present inventive methods and compositions. In this regard, it is expected that amounts
of between 2 mg/kg to about 300 mg/kg will provide fever-reducing therapeutic

SUBSTITUTE SHEET (RULE 26)

activity. Of course, the amount/dose used will depend in specific cases on the particular pharmacological characteristics of the NSAID or combination of NSAIDs included. Further defined ranges of the drug expected to provide the anti-pyretic activity herein disclosed range from between about 10 to about 150 mg/kg or about 20 or 50 mg/kg to about 150 mg/kg.

The present inventors have also observed the claimed compositions are useful for enhancing the platelet retarding activity of a non-steroidal anti-inflammatory drug. This method for inhibiting platelet aggregation comprises providing a non-covalently associated combination of zwitterionic phospholipid and an amount of non-steroidal anti-inflammatory agent that provides reduced inhibition of platelet aggregation in the absence of zwitterionic phospholipid. In some embodiments, this composition is essentially free of anionic phospholipid and/or includes an amount of anionic phospholipid that is a biologically inert component of the preparation. The amounts of non-steroidal anti-inflammatory drug employed as part of the composition, again, is relatively low, and may be further described as an amount that generally provides reduced pharmacological activity in the absence of zwitterionic phospholipid. Again, the zwitterionic phospholipid of choice is, in some embodiments, DPPC, PC, or a combination thereof.

The compositions of the aforescribed methods may further include a neutral lipid, such as a triglyceride. For a partial listing of representative neutral lipids, such as the triglycerides, reference is specifically made to U.S. Patent Number 4,950,656

and 5,043,329. Both saturated and unsaturated triglycerides may be employed in the present compositions, and include such triglycerides as tripalmitin (saturated), triolein and trilinolein (unsaturated). However, these particular triglycerides are listed here for convenience only, and are merely representative of a variety of useful triglycerides, and is further not intended to be inclusive.

Turning now to another aspect of the invention, methods for enhancing the analgesic activity of a non-steroidal anti-inflammatory drug are provided. These methods again comprise providing a non-covalently associated composition comprising zwitterionic phospholipid and an amount of a non-steroidal anti-inflammatory drug that provides reduced pharmacological activity in the absence of zwitterionic phospholipid. In some embodiments, these compositions are essentially free of anionic phospholipid, or include amounts of anionic phospholipid that are biologically inert. In particular embodiments, the non-steroidal anti-inflammatory drug is one or more of those listed in Table 1. In particular embodiments, the NSAID is aspirin, salicylate, a salt thereof, or a combination thereof. While any of a number of different zwitterionic phospholipids may be employed in the composition of the method, in some embodiments, the phospholipid is dipalmitoyl phosphatidyl choline, phosphatidyl choline, or a combination thereof. In these and other embodiments of the described method, an anionic phospholipid that may be excluded or included only in biologically inert amounts is phosphatidyl glycerol (PG).

In still another aspect, the invention provides methods of enhancing the anti-inflammatory activity of non-steroidal anti-inflammatory drugs. The method comprises providing a non-covalently associated combination of zwitterionic phospholipid with an amount of a non-steroidal anti-inflammatory drug. The amount of NSAID in the composition is again defined as an amount that provides reduced pharmacological activity in the absence of zwitterionic phospholipid. In some embodiments, the method employs compositions that are essentially free of anionic phospholipids, such as the DPPG or PG, or includes amounts of anionic phospholipid that are biologically and pharmacologically inert.

In particular embodiments, the composition is further defined as comprising an equimolar amount of NSAID and zwitterionic phospholipid. The present inventors' studies demonstrate the described combination of ingredients provides an enhancement of the anti-inflammatory activity and potency of these drugs compared to drug preparations that do not include zwitterionic phospholipid. Also demonstrated is the reduction in anti-inflammatory activity observed where pharmacologically active (i.e., non-biologically insert amounts) amounts of anionic phospholipid, such as PG (PI), are included in the preparation. Hence, preparations that include pharmacologically active amounts of DPPG or PI would not necessarily provide for the same enhanced anti-inflammatory activity, or the aforescribed enhanced pharmacological activity (i.e., antipyretic or enhanced reduction in platelet aggregation activity), described in the present methods. Pharmacologically active amounts of other negatively-charged

phospholipids would also not be contemplated as particularly useful in view of these results.

Turning now to still a further aspect of the present invention, a method for enhancing the antipyretic potential of subtherapeutically effective amounts of nonsteroidal anti-inflammatory drug is disclosed. In some embodiments, the method comprises again combining zwitterionic phospholipid with an amount of non-steroidal anti-inflammatory drug to provide a noncovalently associated composition thereof. In some embodiments, the composition is further defined as essentially free of biologically active amounts of anionic (or negatively charged) phospholipid.

As used in the description of the above method, a sub-therapeutically effective amount of NSAID is defined as an amount that provides reduced antipyretic activity in the absence of a zwitterionic phospholipid. The enhancement in activity of low amounts of NSAID illustrated in the various *in vivo* studies disclosed herein demonstrate that while doses of aspirin of about 9 mg/kg in combination with the zwitterionic phospholipid, DPPC, did provide for a fever reducing (anti-pyretic) pharmacologic response, treatment with this same dose of NSAID without phospholipid did not.

The present invention also discloses particular pharmaceutical preparations. These pharmaceutical preparations are further described as suitable for enteral or oral administration, and comprise a non-covalently associated combination of zwitterionic phospholipid, non-steroidal anti-inflammatory drug, and a pharmaceutically acceptable

carrier. These compositions are formulated to provide a non-covalently linked composition that is further described as being essentially free of biologically active (pharmacologically active) amounts of anionic phospholipid, and in some embodiments, essentially free of DPPG.

5

In some applications, the non-steroidal anti-inflammatory drug is one or more of those listed in Table 1, such as naproxen, indomethacin, diclophenac, salicylate, aspirin, or any mixture thereof. In particular embodiments, the non-steroidal anti-inflammatory drug of choice is salicylate. While any variety of zwitterionic phospholipids may be employed alone or in combination with the described drugs, some of the representative phospholipids include phosphatidyl choline, dipalmitoyl phosphatidylcholine, phosphatidyl serine, other zwitterionic phospholipids, or mixtures thereof. The compositions may further include a neutral lipid, such as a triglyceride. Representative triglycerides are described in U.S. Patent 4,950,656, which reference is specifically incorporated herein by reference for this purpose. In particular embodiments, the pharmaceutical preparation is defined as comprising an equimolar amount of zwitterionic phospholipid and NSAID.

10

15

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, anti-oxidant, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions and

20

SUBSTITUTE SHEET (RULE 26)

methods described herein is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

As used in the description of the present invention, the term "sub-therapeutically effective amount", particularly as it is used to describe the amount of NSAID employed, is defined as an amount of the NSAID that provides reduced pharmacological (i.e., anti-pyretic) activity in the absence of non-covalent association with a zwitterionic phospholipid.

It is understood that as used in the present disclosure and appended claims, the terms "a" and "an," as in "an element" or "a molecule" are intended to include one or more items or elements, and in no way limit the description or claimed element to one element or item.

The following abbreviations are employed in the description of the invention:

PI = phosphatidyl inositol

PC = phosphatidylcholine

PG = phosphatidylglycerol

L-NAME = N-nitro-L-arginine Methyl Ester

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1A. Photograph of test tubes containing 30 nM of the sodium salts of one of the five NSAIDs tested in chloroform in the absence and presence of 30 nM DPPC (naproxen, indomethacin, diclofenac, salicylate, aspirin).

FIG. 1B. The concentration of each NSAID in solution increases (quantified by UV absorption) in proportion to the molar equivalents of DPPC dissolved in chloroform (Naproxen = -▲-; Aspirin = -●-; Salicylate = -□-, Diclofenac = -△-; Indomethacin = -○-).

FIG. 2A. The DPPC-induced reduction in ASA's solubility in water was maximal when both reactants were present in equimolar concentration over a 60 min. incubation period at 25°C.

FIG. 2B. Time course at 25°C of the reduction in ASA's solubility in water induced by the presence of equimolar concentrations of DPPC. In contrast, the concentration of ASA in water was not significantly changed over time by the addition of an

equimolar concentration of the anionic phospholipid, DPPG as a lipidic suspension

(ASA only = \bullet ; DPPG/ASA = \square ; DPPC/ASA = \circ).

FIG. 3. The passive diffusion of ASA from water into cyclohexane is markedly
5 accelerated by the presence of zwitterionic (DPPC) but not anionic (DPPG)
phospholipids in the aqueous solution (ASA/DPPC = \circ ; ASA/DPPG = \square ; DPPC
= \bullet ; ASA = x).

FIG. 4A. The injurious potential of salicylate and ASA to induce gastric lesions (in
10 rats subsequently challenged with 0.6N HCl), is remarkably decreased when the
NSAIDs are intragastrically administered in association with DPPC (without DPPC =
clear bar; with DPPC = hatched bar).

FIG. 4B. The injurious potential of non-salicylate NSAIDs to induce GI bleeding (in
15 rats pre- and post-treated with L-NAME to increase the animal's susceptibility to the
NSAID) is remarkably decreased when the drugs are intragastrically administered in
association with DPPC (without DPPC = clear bar; with DPPC = hatched bar).

FIG. 4C. The anti-pyretic activity of the NSAID (aspirin - dose 90 mg/kg) is not
20 diminished, and is augmented if administered as a lipidic suspension. Asterisk (*)
represents a statistically significant difference between the values of rats treated with
the NSAID alone (-DPPC) and those treated with the NSAID/DPPC complex (+
DPPC) (Saline = \bullet ; ASA = \square ; ASA/DPPC = \circ).

FIG. 5. Anti-inflammatory action of ASA and Phospholipon G™ (Phospholipid "G") as determined by the implanted string assay as described. Dosage is 90 mg/kg ASA.

Each bar represents data for n=5 rats.

5 FIG. 6. Antipyretic activity of ASA\DPPC complex at a subthreshold ASA dosage (9.0 mg/kg) (Saline = -□-; ASA (9.0 mg/kg) = -▲-; ASA/DPPC (9 mg/kg) = -○-).

FIG. 7. Antipyretic activity of aspirin (ASA) alone when administered at doses which range from 2.5-90.0 mg/kg. In this and all subsequently figures the test agents were
10 intragastrically administered 18 hrs after the rats were subcutaneously injected with 2g/kg Brewer's Yeast to induce a 0.5-1.0°C increase in body temperature. It can be appreciated that doses of aspirin of < 10 mg/kg failed to reduce the fever during the 4 hour study period (Saline = -◇-; ASA, 2.5 mg/kg = -□-; ASA, 5 mg/kg = -○-; ASA, 10 mg/kg = -△-; ASA, 20 mg/kg = -■-; ASA, 45 mg/kg = -●-; ASA, 90
15 mg/kg = -▲-).

FIG. 8. In contrast to the above pattern aspirin (ASA) when complexed with an equimolar concentration of dipalmitoylphosphatidylcholine (DPPC) effectively reduced fever at a dose of 9.0 mg/kg, whereas the anionic phospholipid, DPPG, failed to
20 augment aspirin anti-pyretic activity at this same sub-threshold dose. This figure also demonstrates that anionic phospholipid is essentially biologically inert in terms of enhancing the anti-pyretic action of aspirin (ASA, 9.0 mg/kg = -●-; ASA/DPPC, 9.0

SUBSTITUTE SHEET (RULE 26)

mg/kg = \square -; ASA/DPPG, 9.0 mg/kg = \circ -; n=5/grp; * = p < 0.05 vs ASA; † = p < 0.05 vs ASA/DPPG).

FIG. 9. ASA at a subthreshold dose of 4.5 mg/kg, which had no antipyretic activity alone, effectively reduced fever when complexed with an equimolar concentration of DPPC (H_2O = \square -; H_2O /DPPC (4.5 mg/kg) = Δ -; ASA (4.5 mg/kg) = \ominus -; ASA/DPPC (4.5 mg/kg) = \circ -).

FIG. 10. Compilation of all the data at subthreshold doses (2.5-9.0 mg/kg) of ASA. ASA alone (open symbols) data demonstrates the enhancement of antipyretic activity when ASA was complexed to DPPC (closed symbols) (Saline = $*$ -; ASA, 9mM = \square -; ASA, 4.5 mM = \circ -; ASA, 2.5 mM = Δ -; ASA/DPPC, 9mM = \blacksquare -; ASA/DPPC, 4.5mM = \bullet -; ASA/DPPC, 2.25 mM = \blacktriangle -).

FIG. 11. Dose-response analysis of the antipyretic activity of ASA alone and the ASA/DPPC (equimolar ratio) complex 1 hr (FIG. 11) (ASA = \bullet -; ASA/DDPC = \square -) after intragastric administration. The potency of the ASA, as reflected by the ED_{50} is increased ~ 10 fold when it is administered with the zwitterionic phospholipid.

FIG. 12. Dose response analysis of the anti-pyretic activity of ASA alone and the ASA/DPPC (equimolar ratio) complex 2 hours after intragastric administration.

SUBSTITUTE SHEET (RULE 26)

FIG. 13. Effect of varying the ASA:DPPC ratio from 1:1 on antipyretic activity at 1 hr post intragastric administration. It can be appreciated that the ability of the zwitterionic phospholipid to enhance the antipyretic activity of ASA was lost when the molar concentration of DPPC was increased (from unity) by a factor of 4 or decreased by a factor of 10 (ASA = 9.01 mg/kg).

FIG. 14. Effect of varying the ASA:DPPC ratio from 1:1 on the antipyretic activity of the complex 2 hours post intragastric administration (Legends same as in FIG. 13) (ASA = 9.01 mg/kg).

FIG. 15. At subthreshold doses the antipyretic efficacy of ASA could be clearly enhanced (0-120 min. post-administration), if the NSAID was administered as a microemulsion containing DPPC and tripalmitin (TP). In all cases the molar ratio of ASA:DPPC was maintained at 1:1, whereas the TP was administered in excess (weight ratio of DPPC:TP = 1:4) (n = 5/grp; * = p > 0.05 vs ASA/DPPC; ASA/DPPC 1 mg/kg = -●-; ASA/DPPC/TP, 1 mg/kg = --○--; ASA/DPPC, 9 mg/kg = --▲--; ASA/DPPC/TP, 9 mg/kg = --△--).

FIG. 16. It can be appreciated that the addition of the neutral lipid provided a further enhancement of antipyretic activity, even over that of ASA/DPPC, as can best be seen with the 1 mg/kg subthreshold ASA dose (n = 10/grp; * = p < 0.05 vs ASA/DPPC; ASA, 1 mg/kg = -◆-; ASA/DPPC = -○-; ASA/DPPC/TP = --●--).

FIG. 17. The ability of DPPC to promote the movement of aspirin from water into a lipidic cyclohexane phase (as a membrane model) is greatly accelerated by the presence of the neutral lipid, tripalmitin (TP) to form a microemulsion. This may provide an explanation why the presence of neutral lipids further promote the therapeutic potency of the NSAID/DPPC complex (DPPC + TP (1:4), 2.5mM + ASA, 10mM (sonicated) = -o-; DPPC, 2.5mM + ASA, 10mM (sonicated) = -□-; TP, 2.5 mM + ASA, 10 mM = -■-).

FIG. 18. The ability of DPPC to enhance the anti-pyretic activity of ASA was seen with indomethacin (Saline = -●-; indomethacin (10 mg/kg) = -□-; indomethacin (10 mg/kg)/DPPC = -o-).

FIG. 19. Demonstrates the ability of DPPC to enhance the anti-pyretic activity of naproxen (25 mg/kg dose) (Saline = -●-; naproxen (25 mg/kg) = -□-; naproxen (25 mg/kg)/DPPC = -o-).

FIG. 20. Demonstrates the ability of DPPC to enhance the anti-pyretic activity of diclofenac, 10 mg/kg (Saline = -□-; diclofenac (10 mg/kg) = -□-; diclofenac (10 mg/kg)/DPPC = -o-; n = 5/grp; * = p<0.05 vs. DICLO).

FIG. 21. Demonstrates the ability of DPPC to enhance the anti-pyretic activity of salicylic acid (SA) (70 mg/kg dose) (Saline = -◆-; SA (70 mg/kg) = -o-; SA (70 mg/kg)/DPPC = -●-; n = 5/grp, * = p<0.05 vs. SA/DPPC).

FIG. 22. Antipyretic activity of ASA (18 mg/kg) and ASA/DPPC complex with IP injection of omeprazole (150 mg/kg) two hours prior to NSAID dosage. ASA (saline) = - ● -; ASA/DPPC (Sal) = - ○ -; ASA (omeprazole) = --▲--; ASA/DPPC (omep) = --△--; n = 5/grp; * = p < 0.05 vs. ASA (omep); † = p < 0.05 vs. ASA (Sal).

5

FIG. 23. Antipyretic activity of ASA (18 mg/kg) and ASA/DPPC complex with IP injection of Ranitidine (2 mg/kg) 1 hour prior to NSAID dosage. ASA (saline) = - ● -; ASA (saline) = - ○ -; ASA (Ranitidine) = --▲--; ASA/DPPC (Ranitidine) = --△--; n = 5/grp; * = p < 0.05 vs. ASA (rani); † = p < 0.05 vs. ASA (Sal).

10

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention arises from the discovery that aspirin and other NSAIDs chemically associate with zwitterionic phospholipids, such as dipalmitoyl phosphatidylcholine (DPPC). Important embodiments of the invention include methods of enhancing the various therapeutic activities of NSAID's, such as anti-pyretic, anti-inflammatory, and analgesic pharmacological activities. Surprisingly, these responses are observed without evidence of gastrointestinal side effects as demonstrated in acute and chronic animal models of NSAID injury in the present disclosure.

20

The data disclosed herein indicates that NSAIDs have the capacity to chemically associate with zwitterionic phospholipids in both organic and aqueous solvent systems, and in doing so, both classes of molecules undergo profound changes in their physical and chemical properties. Complex formation in aqueous solvent systems is shown to


occur more efficiently at pH values at or slightly below the pKa of the NSAID.

~~Therefore, without being bound by any theory, it is contemplated that the~~
intermolecular bonding is not covalent, but is instead both hydrophobic and
electrostatic, with the latter association being between the negatively charged carboxyl
5 group of the NSAID and the positively charged nitrogen of the phospholipid. This
possible interaction has been supported by computer assisted molecular modelling
programs (Quanta and CHARMM), which also indicate that the NSAID/phospholipid
complex has a lower molecular free energy (greater thermodynamic stability) than
either reactant alone.

10 According to the present invention, it would be expected that orally
administered NSAIDs would chemically associate with the intrinsic zwitterionic
phospholipids that coat the luminal aspects of the mucus gel layer of the upper GI tract.
A description of luminal aspects of the mucus gel layer is described in Goddard *et al.*,
15 (1990) and Kao *et al.* (1990). While not intending to be limited to any particular
mechanism of action, this intermolecular association is thought to be the basis for the
attenuation in surface activity and/or the loss of stability of the interfacial extracellular
phospholipid layer, and to culminate in an NSAID induced decrease in mucosal
hydrophobicity and barrier properties.

Description of the Lipid Compounds

The phospholipids of the present invention are characterized generally by the formula:



wherein R_1 and R_2 are saturated or unsaturated substitutions ranging from 8 to 32 carbon atoms; R_3 is H or CH_3 , and X is H or COOH ; and R_4 is = O or H_2 .

As will be appreciated by those of skill in the art, the foregoing chemical structure defines a zwitterionic phospholipid structure and embraces a wide range of phospholipids, including but not limited to phosphatidyl cholines, phosphatidyl ethanolamines, phosphatidyl serines and various other zwitterionic phospholipids.

Other phospholipids that may be employed in the composition include: phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine, sphingomyelin, and other ceramides, and mixtures thereof.

SUBSTITUTE SHEET (RULE 26)

Description of the NSAID's


A few of the non-steroidal anti-inflammatory agents that may be employed in the methods and compositions disclosed herein include by way of example: pyrazolones, phenylbutazone (4-butyl-1,2-diphenylpyrazolidine-3,5-dione), and oxyphenbutazone (4-butyl-2-(4-hydroxyphenyl)-1-phenylpyrazolidine-3,5-dione), salicylic acid derivatives such as salicylic acid, salicylic acid amide, acetyl-salicylic acid, benorilate (4-acetamidophenyl-o-acetylsalicylate), and diflunisal (5-(2,4-difluorophenyl)-salicylic acid); Indoles, especially indometacin and its analogs such as indometacin (1-(p-chlorobenzyl)-5-methoxy-2-methylindole-3-acetic acid), glucametacin (1-(p-chlorobenzoyl)-5-methoxy-2-methylindole-3-yl acetic acid glucose amide), acemetacin (1-(p-chlorobenzoyl)-5-methoxy-3-methylindole-3-acetic acid-glycolic acid-ester), and sulindac (5-fluor-2-methyl-1-p-(methylsulphenyl)-benzylidene-indene-3-acetic acid); Phenyl acetic acid or phenyl propionic acid derivatives such as ibuprofen (2-(4-isobutylphenyl)-propionic acid); naproxen (2-(6-methoxy-2-naphthyl)-propionic acid), alclofenac (4-allyloxy-3-chlorophenyl-acetic acid), ketoprofen (2-(3-benzylphenyl)benzoic acid), diclofenac (2-(2,6-dichlorophenylamino)-phenylacetic acid), fenoprofen (2-(3-phenyloxyphenyl)-acetic acid), tolmetin (1-methyl-5-(p-toluy)-pyrrole-2-yl-acetic acid), flurbiprofen (2-(2-fluorobiphenyl-4-yl)-propionic acid), and suprofen (p-2-thenoyl-hydratropic acid) phenyl-propionic acid); Anthranilic acids and their nitrogen analogs such as flufenamic acid (N-(m-trifluoromethylphenyl)-anthranilic acid), mefenamic acid (N-(2,3-dimethylphenyl)-anthranilic acid), and niflumic acid (2-(3-trifluoromethylaminolino)-nicotinic acid).

Phospholipid compounds found to be particularly useful in the practice of the present invention are dilinoleoyl phosphatidylcholine (DLL-PC), dipalmitoyl phosphatidylcholine (DPPC) and egg phosphatidylcholine (Egg-PC or PC_e). In DPPC, a saturated phospholipid, the saturated aliphatic substitution R₁ and R₂ are CH₃ -- (CH₂)₁₄, R₃ is CH₃ and X is H. In DLL-PC, an unsaturated phospholipid, R₁ and R₂ are CH₃ -- (CH₂)₄ -- CH = CH -- CH₂ -- CH = CH -- (CH₂)₇, R₃ is CH₃ and X is H. In Egg PC, which is a mixture of unsaturated phospholipids, R₁ primarily contains a saturated aliphatic substitution (e.g., palmitic or stearic acid), and R₂ is primarily an unsaturated aliphatic substitution (e.g., oleic or arachidonic acid).

Description of the Neutral Lipids

Neutral lipids form another component of some embodiments of the compositions described herein. This class of lipids include the triglycerides.

The triglycerides useful in the practice of the present invention are generally characterized by the formula:



wherein R_1 , R_2 and R_3 are each saturated or unsaturated substitutions ranging from 4 to 32 carbon atoms; and R_4 is either $=O$ or H_2 .

As will be appreciated, this structure embraces a wide range of triglycerides, both saturated and unsaturated, and include, for example, triglycerides such as tripalmitin (saturated), triolein and trilinolein (both unsaturated). A further listing of saturated and unsaturated fatty acids that can be esterified or ether-linked to the triglyceride in question is provided in U.S. 5,032,585, which is specifically incorporated herein by reference.

In a particular anticipated pharmaceutical preparation, the preparation will be provided in a pill form suitable for human ingestion, and contain about 2 to about 300 mg per kg aspirin or salicylate, together with an equimolar amount of PC, DPPC, or a combination thereof, or any other zwitterionic phospholipid. In the described methods, the compositions include the NSAID and the zwitterionic phospholipid in molar ratios ranging from about 1:0.1 to about 1:20, and preferably from about 1:0.5 to about 1:2. In a most preferred embodiments, the ingredients are included in a molar ratio of about 1:1.

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to

constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

5

EXAMPLE 1
GI LEGIONARY AND BLEEDING - EFFECT OF
NSAID AND ZWITTERIONIC PHOSPHOLIPID COMPOSITIONS

10

The present example demonstrates the utility of the present invention for providing compositions that reduce the GI-related side-effects commonly associated with NSAIDS.

15

In order to retard the ability of NSAIDs to interact with the extracellular phospholipid lining of the mucus gel layer, several NSAIDs were preassociated with zwitterionic phospholipids and the effect was determined in various rat ulcer models. The present example shows that ability of the NSAIDs to induce acute and/or chronic GI lesions and bleeding was remarkably decreased when the drugs were administered as a complex with DPPC or related phospholipids. Surprisingly, the anti-pyretic and anti-inflammatory activity of aspirin appeared to be consistently enhanced when associated with zwitterionic phospholipids. This is in contrast to previous side effects of other formulations that reportedly suffer from reduced therapeutic efficacy or onset (Alpsten *et al.*, 1982; Mojaverian *et al.*, 1987).

20

Ulcer Models

Gastric lesions were acutely induced in rats in accordance with the following techniques. For the salicylate-based NSAIDs, fasted male Sprague Dawley rats (150-200g) were intragastrically injected with saline (control), ASA or salicylate, or the drugs preassociated with an equimolar concentration of DPPC (all solutions adjusted to a pH of 3.1). Ten minutes later, the rats were intragastrically challenged with 1 ml of 0.6 N HCl. Gastric lesions were macroscopically scored 60 minutes later in accordance with a previously outlined method (Lichtenberger *et al.*, 1983, incorporated herein by reference).

In order to investigate the effects of non-salicylate NSAIDs to induce GI bleeding, fasted rats were subcutaneously injected with N-nitro-L-arginine Methyl Ester (L-NAME), 1hr. before and 1 and 6 hrs. after intragastrically receiving 1 ml of the NSAIDs; indomethacin, diclofenac, and naproxen administered alone and in association with an equimolar concentration of DPPC. The Nitric Oxide synthesis inhibitor, L-NAME, is administered before and after the NSAID to increase the rat's sensitivity to the drug in accordance with the method of Chen *et. al.*, *Gastroenterology* 104: A53, 1993). Eighteen hours after receiving the NSAID, the distal half of the intestine was excised and flushed with 10 ml of saline. The hemoglobin (Hb) concentration of the intestinal perfusate was measured as an estimate of GI bleeding in accordance with a previously described method (Lichtenberger *et al.*, 1983).

SUBSTITUTE SHEET (RULE 26)

Rats were treated with ASA over a two week period to investigate the chronic effects of NSAID exposure (in the presence and absence of phospholipids) on hematocrit, gastric mucosal hydrophobicity, and granuloma formation. In order to ensure that the rat had an empty stomach prior to receiving the daily intragastric dose of ASA, they were placed on a reverse lighting schedule (9AM/lights off; 5PM/lights on), and were only provided access to chow during the day (dark period). The test solutions (saline, ASA and ASA/Phospholipon 90G complex) were intragastrically administered between 8AM - 9AM daily during the two week study period. In these chronic exposure experiments, ASA was complexed with an equimolar concentration of Phospholipon 90G (purified soya lecithin, prepared and obtained from Nattermann GmbH of Cologne, Germany) instead of DPPC. At the completion of the study period, blood was collected into a capillary tube for the determination of hematocrit and the stomach was excised for contact angle analysis.

One of a number of NSAIDs was intragastrically administered to rats alone or preassociated with an equimolar concentration of DPPC or Phospholipon 90 G (purified soya lecithin, prepared by Nattermann GmbH of Cologne Germany) in several models of acute and chronic injury of the upper GI tract. Two contrasting animal models were employed to determine the ability of DPPC to protect against acute NSAID injury to the GI tract. For salicylate-based NSAIDs, which primarily induce stomach injury, gastric lesions were scored in fasted rats who were initially treated with aspirin or salicylate alone or complexed with DPPC and challenged 10 minutes later with a supra-physiological dose of HCl. For non-salicylate NSAIDs

(indomethacin, diclofenac and naproxen) which predominantly induce injury to the mid-distal regions of the small intestine, the quantity of intraluminal blood was assessed in the distal half of the small intestine of rats who were pre- and post-treated with the Nitric Oxide synthetase inhibitor, L-NAME (N-nitro-L-arginine methyl ester).

5

The results, shown in FIG. 4A and FIG. 4B, respectively, clearly demonstrate (in rats sensitized to the drugs) that the injurious potential of both salicylate-based and non-salicylate NSAIDs to induce acute GI lesions and bleeding is significantly reduced, by > 85%, if the NSAID is preassociated with the zwitterionic phospholipid prior to administration. Similarly, daily administration of aspirin to rats over a 2 week period resulted in a significant fall in both hematocrit and gastric mucosal hydrophobicity, which was prevented in rats that received the aspirin/Phospholipon G complex (Table 2).

10

15

Table 2. Effect of DPPC on Aspirin's Chronic GI Side Effects

Group	Gastric Mucosal Hydrophobicity (degrees, Contact Angle) ^b	Hematocrit
Saline (Control)	52.2 ± 2.7(6)	51.7 ± 0.4 (10)
Aspirin (90mg/kg)	15.8 ± 6.6 ^c (6)	47.8 ± 0.2 ^c (10)
Aspirin + Phospholipon G	55.3 ± 5.8 ^{c,d} (6)	53.1 ± 0.4 ^{c,d} (10)

^b Gastric mucosal hydrophobicity was measured by contact angle analysis as described in Hills *et al.*, 1983; Goddard *et al.*, 1987; Goddard *et al.*, 1990.

^c $p < 0.05$ in comparison to saline-treated control values.

^d $p < 0.05$ in comparison to values of aspirin-treated rats.
(n) number of rats/group

EXAMPLE 2 - SOLUBILITY STUDIES; COMPLEX FORMATION BETWEEN NSAIDS and ZWITTERIONIC PHOSPHOLIPIDS

To demonstrate the effect of DPPC on the solubility of the sodium salts of the five NSAIDs (naproxen, indomethacin, diclofenac, salicylate and aspirin) in chloroform, each NSAID was added to chloroform at a 30 nM final concentration. DPPC was dissolved in the chloroform that was contained in half the tubes at a final concentration which ranged between 5 - 40 nM, prior to the addition of the NSAID salt. DPPC, as well as PC and other zwitterionic phospholipids useful in the practice of the invention, may also be dissolved in other organic solvents, such as ethanol, in the practice of the present invention. The tubes were gently mixed at 25°C for 16 hrs, after which they were photographed and/or centrifuged (2,000g for 15 min) and the supernatant collected to determine the concentration of NSAID

SUBSTITUTE SHEET (RULE 26)

in solution. The latter was assessed by measuring the NSAID's UV absorbance at 290 nm, and comparing these readings to the appropriate standard curves for each

NSAID. It should be noted that for each of the five NSAIDs a linear relationship existed between the drug's concentration (nM) in water and the UV absorbance reading. The equations for the regression lines for each of the NSAIDs were as follows: aspirin, $y = 2.603x - 0.004$, $r = 0.999$; salicylate, $y = 8.780x + 0.123$, $r = 0.999$; indomethacin, $y = 18.325x + 0.156$, $r = 0.999$; diclofenac, $y = 21.523x + 0.008$, $r = 0.999$; and naproxen, $y = 3.732x + 0.005$, $r = 0.999$. In all cases the readings for the unknowns fell within the linear portion of the standard curves. Further, the presence of DPPC in the solvent did not interfere with these analyses, as it failed to contribute to the UV absorbance reading in the absence of the NSAID.

To assess the effect of DPPC on the solubility of the sodium salt of ASA in water, ASA was dissolved in water at a final concentration of 30 nM (pH adjusted to 6.0), and its intrinsic fluorescence (290 nm/excitation; 406 nm/emission) was monitored. DPPC was present as a lipidic suspension in half the tubes at a final concentration which ranged between 15 - 60 nM. The tubes were gently mixed at 25°C for the desired incubation period, after which they were centrifuged (2,000g for 15 min) and the supernatant collected to determine the concentration of ASA in solution. Once again a linear relationship ($y = 14.02 + 0.353x$, $r = 0.999$) was found between the fluorescent reading and the concentration of aspirin in solution.

It can be appreciated from FIG. 1A and FIG. 1B that the sodium salts of these NSAIDs are insoluble in chloroform unless an equimolar or greater

concentration of DPPC is added to the organic solvent, at which point complete solvation takes place. Conversely the sodium salts of the NSAIDs are readily soluble in either saline or water, and are removed from solution as a complex within minutes after an equimolar concentration of DPPC is added as a lipidic suspension. The solubility of aspirin in saline can be followed either by monitoring its intrinsic fluorescence or radioactivity (employing ^{14}C -labelled aspirin). FIG. 2A and FIG. 2B demonstrate that the injection of DPPC into an aqueous solution results in the precipitation of the NSAID, presumably as a complex with the phospholipid. The rapid change in solubility of the NSAID in both the organic and aqueous solvent systems does not occur if DPPC is substituted by the anionic phospholipid, dipalmitoylphosphatidylglycerol (DPPG).

EXAMPLE 3 - GRANULOMA FORMATION AND ENHANCED ANTI-INFLAMMATORY ACTIVITY

The present example demonstrates the utility of the present methods for enhancing the anti-inflammatory activity of ASA accomplished when ASA, or other non-steroidal anti-inflammatory drug complexed with Phospholipon 90G. A foreign-body granuloma model widely used by those of skill in the art to assess anti-inflammatory action was employed.

The inventors used the rat model of foreign-body granuloma formation. This model is recognized by those of skill in the art as a representative model for

granuloma formation and anti-inflammatory activity, as described in Ucelay *et al.*,
(1988); and Castro *et al.*, (1980). These references are specifically incorporated
herein for the purpose of providing details associated with the use of this model.

5

Methods

The above model has been successfully employed in both rats and
invertebrates to quantify this most basic component of a tissue's response to injury
(Ucelay *et al.*, 1988; Castro *et al.*, 1980; Clatworthy *et al.*, 1994, each
incorporated herein by reference). Sterile tared cotton string was surgically
implanted (bilaterally) under the abdominal skin of ether anesthetized rats on day 1
of the study period, and then randomly placing rats in a group to be daily treated
with saline, aspirin or the NSAID complexed with Phospholipon 90G over a two-
week period.

15

At the completion of the study period the string with the adherent granuloma
tissue was surgically dissected from the euthanized rats and dried in a vacuum for
several days at room temperature, until a baseline dry weight was obtained. The
difference between this value and the initial dry weight of the string prior to
implantation divided by the latter value provided an estimate of the weight of
granuloma tissue. This technique proved to be very reproducible and accurate, as
determined both by the close agreement between the changes in weight of the two
pieces of string that were implanted contralaterally in each animal ($< 12.5\%$

20

difference in values between the right and left string); and the low variance ($< 8\%$) in values of granuloma formation within a group of animals.

Improvement in the anti-inflammatory efficacy of ASA was observed over a 2 week study period. This effect is seen when the NSAID was administered as a complex with Phospholipon G at concentrations below the maximally effective dose for this particular drug (see Table 3, FIG. 5).

Table 3. Effect of Aspirin \pm DPPC Formulations on Foreign-Body Granuloma Formation.

Group	\pm DPPC	n	Granuloma Formation ^a
Saline (Control)	-	5	3.20 ± 0.10
ASA (90 mg/kg)	-	5	1.90 ± 0.10^b
	+	5	$1.4 \pm 0.10^{b,c}$
ASA (140 mg/kg)	-	6	1.07 ± 0.09^b
	+	6	1.00 ± 0.15^b

^a The values represent the dry weight of the string with adherent granuloma tissue - dry weight of string/dry weight of the string.

^b $p < 0.05$ in comparison to saline-treated control values.

^c $p < 0.05$ in comparison to values of rats treated with NSAIDs alone.

EXAMPLE 4 - CONTACT ANGLE ANALYSIS

The present example is provided to describe the model which was used to examine the surface tension reducing action of the compositions of a combination of non-steroidal anti-inflammatory agent and zwitterionic phospholipid.

SUBSTITUTE SHEET (RULE 26)

Contact angle analysis was performed with the use of a goniometer on excised gastric mucosal tissue, that was lightly blotted and dried, as previously outlined (Hills *et al.*, 1983; Goddard *et al.*, 1987; Goddard *et al.*, 1990, each incorporated herein by reference). Briefly, this was accomplished by applying a droplet of water ($\sim 5\mu\text{l}$) to the tissue surface, and employing the telescopic eyepiece of the goniometer to measure the maximal angle that is dissected at the triple point, where the solid/liquid/ and air interface meet.

EXAMPLE 5 - ANTIPYRETIC ACTIVITY

An established rat fever model was used in the present example to demonstrate the utility of the invention for enhancing the anti-pyretic activity of a NSAID by combining these class of agents with a phospholipid.

The rat fever model involves the injection of rats with Brewer's yeast (2g/kg, s.c.) to induce an increase in fever of 0.5 -1.5°C. These models are described in Adams *et al.* (1968) and Ucelay *et al.* (1988), which references are specifically incorporated herein for this purpose. The animals were intragastrically treated (instilled) with either saline, 90 mg/kg ASA, or 90 mg/kg ASA preassociated with an equimolar concentration of DPPC. Similar antipyretic analyses were performed with the sodium salts of the following NSAIDs: diclofenac (10 mg/kg), indomethacin (10 mg/kg) and naproxen (30 mg/kg), alone and complexed with an equimolar concentration of DPPC. All test solutions were titrated to a pH of 4.5 prior to intragastric administration. Rectal temperatures were

SUBSTITUTE SHEET (RULE 26)

monitored in conscious, restrained rats at all indicated times. This technique
provided a very reliable and reproducible estimate of the antipyretic activity of

NSAID formulations as indicated by the fact the variance within a group was low,
with the standard errors $< 5\%$ of the mean values, and the fact that the difference
in mean temperature values for a given group varied $< 2\%$ between separate
experiments.

EXAMPLE 6 - PHARMACEUTICAL COMPOSITIONS

The complexes described in the present example are preferably for oral
administration, for example, with an inert diluent or with an assimilable edible
carrier, or they may be enclosed in hard or soft shell gelatin capsule, or they may
be compressed into tablets, or they may be incorporated directly with the food of
the diet. For oral therapeutic administration, the active compounds (i.e., NSAID
and Zwitterionic phospholipid and/or neutral lipid) may be incorporated with
excipients or carriers and used in the form of ingestible tablets, buccal tablets,
troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such
compositions and preparations should contain enough active NSAID compound to
achieve an effective plasma level of the active drug. Aspirin, for example, would
be provided in doses of from about 10 mgm, or about 20 mg, or 32.5 mg or even
up to 60 mg, or 300 mgs/kg would be contained in each tablet or dose, where
employed for administration to animals having a weight of about 60 kg - 70 kg.
The dose will vary depending on the NSAID or combinations of NSAIDs used.
The amount of the NSAID in particular preparations is more generally described as

SUBSTITUTE SHEET (RULE 26)

an amount of NSAID or combination of NSAIDs effective to provide a
~~pharmacologically active plasma concentration of the drug when used in~~
combination with zwitterionic phospholipid. These amounts are determinable by
one of ordinary skill in the pharmaceutical arts given the data disclosed herein, and
5 general pharmaceutical references such as Remingtons Pharmaceutical Sciences.

The tablets, troches, pills, capsules and the like may also contain the
following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients,
such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato
10 starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a
sweetening agent, such as sucrose, lactose, aspartame or saccharin may be added,
or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring.
When the dosage unit form is a capsule, it may contain, in addition to materials of
the above type, a liquid carrier. Various other materials may be present as coatings
15 or to otherwise modify the physical form of the dosage unit. For instance, tablets,
pills, or capsules may be coated with shellac, sugar or both.

A syrup or elixir may contain the active compounds sucrose as a sweetening
agent methyl and propylparabens as preservatives, a dye and flavoring, such as
20 cherry or orange flavor, again formulated for oral administration. Of course, any
material used in preparing any dosage unit form should be pharmaceutically pure
and substantially non-toxic in the amounts employed. In addition, the active

compounds may be incorporated into sustained-release preparation and formulations.

The present compositions may also be formulated as injectable formulations, or as formulations suitable for enteral administration, according to those techniques known to those of ordinary skill in the medicinal arts.

EXAMPLE 7

Lipid-Permeability of NSAID/Phospholipid Complex

The present example illustrates the utility of the invention for providing compositions and methods for enhancing the bio-absorption and bio-availability of NSAID's.

The diffusion of aspirin (alone and complexed with phospholipid) from water into cyclohexane was used as a model to estimate the membrane permeability of the drug.

Permeability Analysis

The sodium salt of ASA (also salicylate) was dissolved in 5 ml of water at a final concentration of 100mM (pH adjusted to 6.0) and gently stirred at 25°C. An equal volume of cyclohexane was layered over the aqueous solution and the entry of the NSAID into the organic phase was monitored fluorometrically over time. In order to determine the effect of phospholipid association on the lipid permeability of the NSAID, ASA (or salicylate) at the above concentration was sonicated in the

presence of 0.5 mM phospholipid (DPPC or DPPG) in water (adjusted to a pH of 7), and its rate of diffusion into the cyclohexane phase was measured

fluorometrically. This was accomplished by removing 1 ml of the top phase chloroform solution by pipette, injecting it into the cuvette to obtain the fluorescence reading, and returning the sample to the incubation vessel to assure that the volume did not change. This entire process could be completed in < 30 seconds.

In these studies, the concentration of NSAID:phospholipid in water was adjusted to a molar ratio of 200:1, creating a large driving force to promote NSAID flux into the hydrocarbon phase, and minimizing the turbidity encountered with high phospholipid concentrations.

Results

Under neutral conditions, the passive diffusion of aspirin across an aqueous - hydrocarbon interface, as assessed fluorometrically, was negligible unless it was chemically associated with the zwitterionic phospholipid, DPPC (See Figure 3). Furthermore, this increase in the flux rate into the organic phase, was simply not a consequence of liposomal encapsulation since the NSAID failed to enter cyclohexane if the anionic phospholipid, DPPG was substituted for DPPC. These studies also revealed that DPPC promoted the flux of sodium-salicylate from the aqueous to the organic phase in a similar manner.

EXAMPLE 8
ANTI-INFLAMMATORY AND ANTI-PYRETIC ACTIVITY
OF NSAID/PHOSPHOLIPID COMPLEX

5 This example demonstrates the utility of the present invention for enhancing
the fever-reducing potential of the NSAID (20 mg/kg dose) when chemically
associated with a zwitterionic phospholipid. The ability of NSAIDs (administered
alone and complexed with DPPC) to reduce fever in rats was determined. Fever
was induced 18 hrs prior to drug treatment by the subcutaneous administration of
10 Brewer's yeast (Adams *et al.*, 1968; Ucelay *et al.*, 1988).

 The results (See FIG. 4C) indicate that the anti-pyretic activity of the
ASA/DPPC complex was significantly greater than that of the NSAID alone, at all
time periods examined.

15 The enhancement in anti-pyretic activity of the NSAID/DPPC complex
compared to the NSAID alone over the first three hours after intragastric
administration, was observed to a lesser degree with sodium salts of the following
drugs; diclofenac (-0.29°C), indomethacin (-0.28°C), and naproxen (-0.30°C).

20

EXAMPLE 9
ENHANCEMENT OF ANTIPYRETIC ACTIVITY OF
ASA/DPPC COMPLEX AT SUBTHRESHOLD ASA DOSAGE

25 The antipyretic activity of the ASA/DPPC complex was determined in rats
as in Example 8 (*supra*), this time at much lower doses, more than 2 times lower
than in Example 8. In the present example, a dosage of 9.0 mg/kg ASA was

SUBSTITUTE SHEET (RULE 26)

administered either alone or complexed with DPPC. The data from this study is summarized in FIG. 6.

As can be seen in FIG. 6, ASA alone does not have significant antipyretic activity at this dosage level, however ASA complexed with DPPC does have significant antipyretic activity over a five hour period. The dosage level in the present example is a 10-fold reduction of the standard dosage of 90 mg/kg, as used in Example 8, and reported in FIG. 4C.

Based on the recommended human dosages of 90 mg/kg for juvenile rheumatoid arthritis, or 325 to 650 mg for antipyretic or analgesic treatment in adults (See pages 1110-1111, Remington's Pharmaceutical Sciences, 18th Edition, Mack Publishing Company, Easton, Pennsylvania, 1990, incorporated herein by reference), it is expected that ASA complexed with DPPC, or other zwitterionic phospholipid, at a 10-fold lower dosage, i.e. approximately 9 mg/kg for juvenile rheumatoid arthritis or 32.5 to 65 mg for antipyretic activity in adults, would be as effective as the normal dosage of ASA alone to provide fever-reducing activity.

EXAMPLE 10 ANALGESIC EFFECT OF COMPLEXED-NSAIDS

It is contemplated that, in light of the demonstrated enhancement of the effects of NSAIDs (anti-inflammatory and antipyretic) when complexed with a zwitterionic phospholipid, or even further administered in combination (i.e., in a mixture) with a neutral lipid (triglyceride) that the analgesic effects of these drugs

will also be enhanced when complexed with the phospholipids. Two established pain tests, using rats, may be employed to demonstrate enhancement of this effect.

Tail retraction occurs in response to a noxious stimulus that generates pain. Tail retraction response may be observed and timed in rats given a placebo in order to determine a base line reaction time. If after receiving either DPPC-complexed or uncomplexed aspirin the time latency between pain induction (with a laser heat source) and tail retraction increases, this is a direct reflection of the analgesia.

Retraction times will be compared between placebo and aspirin-treated rats. The test will be conducted with varying dosages and varying times after the administration of the placebo, aspirin, and/or other NSAIDs, or complex to determine if dosage or duration have some effect on the strength of the analgesia.

A model for analgesic effect in rats involves animals injected with formalin.

The rats are injected under the dorsal surface of the right hind paw with a 0.05 ml volume of 15% formalin and saline solution (Helmstetter and Fanselow, 1987).

The treated rats are then given the aspirin, and/or other NSAIDs, aspirin/DPPC or PC complex, or placebo at various times before and after injection with the irritating solution. The rats are then placed in a cage and their behavioral responses to the painful stimulus are observed (employing a video camera system) and graded as follows: (1) freezing - an absence of all activity other than respiration; (2) paw lifting - the rat holds its treated paw close to its body; (3) paw licking - the rat either licks the treated paw or has some other type of mouth contact with it; and (4) general activity - this involves any other type of general movement. The

SUBSTITUTE SHEET (RULE 26)

behavioral response of the rat is indicative of its sensitivity. The formalin test, along with the tail flick test, comprises two distinct and separate tests to evaluate pain sensitivity and analgesia effectiveness.

5

EXAMPLE 11
Effects of Low Dose Aspirin (9 mg/kg) Alone
and as a Complex with Phospholipid/Neutral Lipids
on Foreign-Body Granuloma Formation in Rats*

The present example is provided to demonstrate the utility of the present invention for reducing inflammation. The string granuloma model described herein was again used to demonstrate the activity of the presently disclosed methods for treating this condition.

Sterile tared string was surgically implanted. Five days later rats were intragastrically administered twice a day with either saline, ASA alone, or ASA lipid mixtures. Rats were sacrificed after each rat received 5 doses of the test compounds and the string and granuloma excised and weighed.

* $p < 0.05$ vs. ASA alone. ASA = Aspirin;

DPPC = Dipalmitoylphosphatidylcholine,

20

TP = Tripalmitin; TO = Triolein

TABLE 3

Saline	ASA	ASA DPPC/TP	ASA PhospholiponG/TO
4.08(8)	3.89(8)	3.44(8)	3.49(8)*
± 25	± 0.06	± 0.24	± 0.17

Table 3. As demonstrated in the data provided in Table 3, a phospholipid (DPPC)/neutral lipid (TO) microemulsion appeared to modestly enhance the anti-inflammatory activity of ASA when the NSAID was administered at a subthreshold dose (9 mg/kg) for this activity.

EXAMPLE 12
Effect of DPPC on Aspirin's Inhibitory
Effect on Platelet Aggregation

The present example demonstrates the ability of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) to inhibit cellular activation resulting in either platelet aggregation or the synthesis and release of inflammatory mediators. The activity is shown to be enhanced if the NSAIDs are administered as a complex with zwitterionic phospholipids alone or together with neutral lipids.

High Ristocetin (1.5 mg/ml) Test		
	% Platelet Aggregation	
ASA Conc	-DPPC	+DPPC
0.00	93%	100%
0.01mM	100%	100%
0.1 mM	83%	87%
1.0 mM	95%	35%
1.0 mM ^a	100%	84%
Low Ristocetin (0.75) mg/ml Test		
	% Platelet Aggregation	
ASA Conc	-DPPC	+DPPC
0.00	75%	94%
0.01mM	65%	12%
0.1mM	26%	20%
1.0mM	80%	6%
1.0mM ^a	39%	15%

Abbreviations: ASA = aspirin

DPPC = dipalmitoylphosphatidylcholine

DPPC was added at a conc. equimolar to the aspirin.

In the absence of aspirin, DPPC was added at a final concentration of 1 mM.

^a Second study.

SUBSTITUTE SHEET (RULE 26)

The advantage of this invention is it will allow the NSAIDs to be administered at a lower than normal dose, due to their enhanced efficacy and potency-thus increasing their effectiveness and minimizing their side-effects on the GI tract and other organ systems.

5

A number of embodiments of the invention would include the combination of NSAIDs with zwitterionic phospholipids alone and together with neutral lipids. These combinations would both increase the efficacy and potency of NSAIDs to inhibit the activation of: 1) platelets; 2) neutrophils; 3) monocytes/macrophages; 4) lymphocytes; 5) Pans and 7) other bone-marrow derived cell types.

10

Considering the interest in the pharmaceutical industry in the role of NSAIDs in the prevention of cardiovascular disease and tissue/joint inflammation, the presently described pharmaceutical preparations would provide alternative clinical management protocols with a improved bioavailability at lower doses of the sometimes irritative NSAID regimen.

15

This list would include at minimum the 20-40 pharmaceutical companies presently marketing an NSAID.

20

EXAMPLE 13**Anti-Secretory Agents together with NSAID and Lipids**

The present example demonstrates the utility of the present invention for compositions that include an anti-secretory drug, such as Tagamet, or other histamine type 2 receptor antagonist, or Omeprazole (Prilosec™, or other proton-pump inhibitor or H^+/K^+ ATPase inhibitors), either before or along with an NSAID complexed with phospholipid and/or neutral lipid (such as a triglyceride). These embodiments of the invention are further described in Example 14.

While not intending to be limited to any particular mechanism of action, by including a phospholipid and/or neutral lipid, the poor absorption of NSAID's that sometimes results with an anti-secretory agent administered therewith, may to some degree be prevented or lessened. Hence, the bioavailability and therapeutic action of the NSAID when administered together with an anti-secretory agent may be maintained, and in some cases enhanced.

The present example also demonstrates that the therapeutic (antipyretic) activity of aspirin, and other NSAIDs, is attenuated if animals are pre-treated with an agent that inhibits gastric acid secretion. Omeprazole (sold under the name Prilosec™) is in the class of "proton pump inhibitors", also called " H^+/K^+ ATPase inhibitors" that act by irreversibly binding to and inhibiting H^+/K^+ ATPase of the perietal cell, the rate limiting enzyme in gastric HCl secretion. Ranitidine (sole under the name Zantac™) is in the class of " H_2 receptor antagonists" that prevents

histamine from binding to its type-2 receptor on the parietal cell to inhibit gastric acid secretion. It can be appreciated from figures 22 and 23 that the blocking effect

of these two classes of antisecretory drugs on the therapeutic actions of the NSAID, however, is overcome if the NSAID is complexed with a zwitterionic phospholipid.

5 As demonstrated in Figures 23 and 23, aspirin at a dose of 20 mg/kg failed to reduce fever in rats if the NSAID was administered in conjunction with either the proton pump inhibitor (Prilosec™) or a histamine blocker, Ranitidine (Zantac™). This block in therapeutic activity due to inhibition of gastric acid secretin was overcome if the NSAID was administered as a complex with phospholipid, DPPC.

10

EXAMPLE 14 - Therapeutic Regimens

The present example is provided to demonstrate various therapeutic combination regimens for treating fever, inflammation and pain. These regimens include the administration of NSAIDs together with phospholipid and/or neutral
15 lipid.

Agents that include phospholipid, such as lecithin - tablets and the like, may be used as part of the regimens disclosed herein for the enhancement of NSAID activity.

20

Therapeutic Regimen

For use as an improved regimen (e.g., anti-pyretic, platelet aggregation, analgesic), the present invention contemplates an initial administration of the

phospholipid, such as in a tablet or phospholipid-containing agent, such as lecithin tablets, that contain phospholipid (e.g., phosphatidyl choline).

5 Either at the same time or following the administration of a phospholipid or phospholipid containing agent, the patient would then be given an NSAID. The NSAID may also be administered in combination with the phospholipid as a single composition, or alternatively as a combination with both a phospholipid and a neutral lipid, such as a triglyceride (e.g., TO).

10 The present inventors also propose regimens that include the administration of an NSAID and phospholipid and/or neutral lipid, either before or at the same time as an anti-secretory drug, such as Tagamet® and Prilosec™. This is because antisecretory drugs are observed by the present inventor to reduce the anti-pyretic action of NSAID's. It is expected that the inclusion of phospholipid and/or neutral
15 lipid will improve the observed reduced absorption of NSAID's observed when NSAIDs are administered with an antisecretory agent alone. Typically, this reduced absorption required that a higher dose of the NSAID be administered to the patient in order to provide the desired therapeutic effect.

20 Combinations of NSAIDs with zwitterionic phospholipids alone or in combination with neutral lipids will promote the ability of this family of drugs to influence certain target cells, such as neutrophils, platelets, eosinophils, macrophages, and others. In doing so, said phospholipids will increase the efficacy

and potency of the NSAIDs to inhibit the cellular cyclo-oxygenase and the formation of arachidonic acid-derived products and other agents involved in cellular aggregation, adhesion, and the synthesis and release of inflammatory mediators and/or cytokines.

5

* * *

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure.

10

While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent

15

that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

20

SUBSTITUTE SHEET (RULE 26)

REFERENCES

The following references, to the extent that they provide exemplary

procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5 Adams, S.S., Hebborn, P. & Nicholson, J.S. Some aspects of the pharmacology of ibufenac, a non-steroidal anti-inflammatory agent. *J. Pharm. Pharmac.* **20**, 305-312 (1968).
- 10 Alexander, A.M., Veitch, G.B.A. & Wood, J.B. Anti-rheumatic and analgesic drug usage and acute gastro-intestinal bleeding in elderly patients. *J. Clin. Hosp. Pharm.* **10**, 89-93 (1985).
- 15 Allison, M.C., Howatson, A.G., Torrance, C.J., Lee, F.D. & Russell, R.I. Gastrointestinal damage associated with the use of nonsteroidal anti-inflammatory drugs *N. Engl. J. Med.* **32**, 749-754 (1992).
- 20 Alpsten, M., Bogentoft, C., Ekenved, G., & Solvell, L. Gastric emptying and absorption of acetylsalicylic acid administered as enteric-coated microgranules. *Eur. J. Clin. Pharm.* **22**, 57-61, 1982.
- Castro, G.A., Malone, C. & Smith, S. Systemic anti-inflammatory effect associated with enteric trichinellosis in the rat. *J. Parasitol.* **66**, 407-412 (1980).
- 25 Fuster, V., Dyken, M.L., Vokonas, P.S., and Hennekens, C. Aspirin as a therapeutic agent in cardiovascular disease. *Circulation* **87**, 659-675 (1993).
- Ghyczy *et al.*, US Patent 4,309,420.
- 30 Ghyczy *et al.*, US Patent 4,369,182.
- Ghyczy *et al.*, US Patent 4,421,747.
- 35 Goddard, P.J., Kao, Y-C. & Lichtenberger, L.M. Luminal surface hydrophobicity of canine gastric mucosa is dependent on a surface mucous gel. *Gastroenterology* **98**, 361-370 (1990).
- 40 Goddard, P.J., Hills, B.A. & Lichtenberger, L.M. Does aspirin damage the canine gastric mucosa by reducing its surface hydrophobicity? *Am. J. Physiol.* **252**, G-421-430 (1987).

Graham, D.Y. Prevention of gastroduodenal injury induced by chronic nonsteroidal anti-inflammatory drug therapy. *Gastroenterology* 96, 675-681 (1989).

5

Helmstetter, F.J., Fanselow, M.S. Strain differences in reversal of conditional analgesia by opiod antagonists. *Behavioral Neurosciences*, 101, 735-737 (1987).

10

Hills, B.A., Butler, B.D. & Lichtenberger, L.M. Gastric Mucosal Barrier: The hydrophobic lining to the lumen of the stomach. *Am. J. Physiol.* 244, G-561-568 (1983).

Jolobe, O.M.P. & Montgomery, R.D. Changing clinical pattern of gastric ulcer: are anti-inflammatory drugs involved? *Digestion* 29, 164-170 (1984).

15

Kao, Y-C., & Lichtenberger, L.M. Phospholipid- and neutral- lipid containing organelles of rat gastroduodenal mucous cells. *Gastroenterology* 101, 7-21 (1991).

20

Kao, Y-C., Goddard, P.J. & Lichtenberger, L.M. Morphological effects of aspirin and prostaglandin on the canine gastric mucosal surface: analysis with a phospholipid cytochemical stain. *Gastroenterology* 98, 592-606 (1990).

Kao, Y-C., & Lichtenberger, L.M. Effect of 16, 16 dimethyl prostaglandin E₂ on the lipidic organelles of rat gastric surface mucous cells. *Gastroenterology* 104, 103-113 (1993).

25

Lichtenberger, L.M., Graziani, L.A., Dial, E.J., Butler, B.D. & Hills, B.A. Role of surface-active phospholipids in gastric cytoprotection. *Science* 219, 1327-1329 (1983)

30

Ligumsky, M., Grossman, M.I. & Kauffman, Jr., G.L. Endogenous gastric mucosal prostaglandins: their role in mucosal integrity. *Am. J. Physiol.* 242, G337-341 (1982).

35

McCormack, K. & Brune, K. Classical absorption theory and the development of gastric mucosal damage associated with non-steroidal anti-inflammatory drugs. *Arch. Toxicol.* 60, 261-269 (1987).

40

Mojaverian, P., Rocci, Jr., M.L., Conner, D.P., Abrams, W.B. & Vlasses, P.H. Effect of food on the absorption of enteric-coated aspirin: Correlation with gastric residence time. *Clin. Pharm. Ther.* 41, 11-17 (1987).

U.S. Patent # 4,369,182

U.S. Patent # 4,421,747

45

U.S. Patent # 4,309,402

Rainsford, K.D. *Antiinflammatory and Anti-rheumatic Drugs I*, CRC Press, Boca Raton, Florida, (1985).

5 Rainsford, K.D. Mechanism of gastrointestinal toxicity of non-steroidal anti-inflammatory drugs. *Scand. J. Gastroenterol.* **24** (suppl. 163), 9-16 (1989).

Ucelay, M., Lasheras, B. & Cenarruzabeitia, E. Pharmacological study of the new nonsteroidal antiinflammatory agent, 4'-Acetamidophenyl-2-(5'-p-toluy-1'-methylpyrrole) acetate. *Arzheim-Forsch/Drug Research* **38**, 546-551 (1988).

10 Whittle, B.J.R. Temporal relationship between cyclooxygenase inhibition, as measured by prostacyclin biosynthesis and the gastrointestinal damage induced by indomethacin in the rat. *Gastroenterology* **80**, 94-98 (1981).

15 Whittle, B.J.R., Higgs, G.A., Eakins, K.E., Moncada, S. & Vane, J.R. Selective inhibition of prostaglandin production in inflammatory exudates and gastric mucosa. *Nature* **284**, 271-273 (1980).

20 Willard, J.E., Lange, R.A. and Hillis, L.D. The use of aspirin in ischemic heart disease. *N. Engl. J. Med.* **327**, 175-181 (1992).

WHAT IS CLAIMED IS:

1. ~~A method for enhancing the anti-pyretic activity of a non-steroidal anti-inflammatory drug comprising:~~

5

providing a non-covalently associated composition of a zwitterionic phospholipid with an amount of non-steroidal anti-inflammatory agent that provides reduced anti-pyretic activity in the absence of the zwitterionic phospholipid.

10

2. The method claim 1 wherein the composition is further defined as comprising an equimolar amount of zwitterionic phospholipid and non-steroidal anti-inflammatory agent.

15

3. The method of claim 1 wherein the non-steroidal anti-inflammatory drug is salicylate, aspirin, naproxen, indomethacin, diclofenac, or a mixture thereof.

20

4. The method of claim 1 wherein the anti-pyretic activity of the non-steroidal anti-inflammatory drug is enhanced about 2-fold to about 6-fold in the presence of the zwitterionic phospholipid compared to non-steroidal anti-inflammatory drug without phospholipid.

25

5. The method of claim 1 wherein the non-steroidal anti-inflammatory drug is salicylate.

30

6. The method of claim 1 wherein the zwitterionic phospholipid is dipalmitoyl phosphatidylcholine.

5 7. The method of claim 1 wherein the amount of non-steroidal anti-inflammatory drug is about 2 mg to about 300 mg.

10 8. A method of inhibiting platelet aggregation comprising:
providing a non-covalently associated combination of zwitterionic
phospholipid and an amount of non-steroidal anti-inflammatory agent
that provides reduced inhibition of platelet aggregation in the absence
of zwitterionic phospholipid.

15 9. The method of claim 8 wherein the composition is further defined as
comprising an equimolar amount of zwitterionic phospholipid and non-steroidal
anti-inflammatory agent.

20 10. The method of claim 8 wherein the zwitterionic phospholipid is dipalmitoyl
phosphatidyl choline.

25 11. The method of claim 8 wherein the non-steroidal anti-inflammatory drug is
salicylate, aspirin, naproxen, indomethacin, diclofenac, or a mixture thereof.

30 12. The method of claim 1 or 8 wherein the composition further comprises
neutral lipid.

SUBSTITUTE SHEET (RULE 26)

13. The method of claim 12 wherein the neutral lipid is triglyceride.

5 14. A method for enhancing the analgesic activity of a non-steroidal anti-inflammatory drug comprising:

10 providing a non-covalently associated composition of a zwitterionic phospholipid and an amount of a non-steroidal anti-inflammatory drug that provides reduced pharmacological activity in the absence of zwitterionic phospholipid.

15 15. A method for enhancing the anti-inflammatory activity of a non-steroidal anti-inflammatory drug comprising:

20 providing a non-covalently associated composition of a zwitterionic phospholipid with an amount of a non-steroidal anti-inflammatory drug that provides reduced pharmacological activity in the absence of zwitterionic phospholipid.

25 16. The method of claim 15 wherein the composition is further defined as comprising an equimolar amount of zwitterionic phospholipid and non-steroidal anti-inflammatory drug.

30 17. The method of claim 14 or 15 wherein the non-steroidal anti-inflammatory drug is salicylate, naproxen, indomethacin, diclofenac, aspirin, or a mixture thereof.

SUBSTITUTE SHEET (RULE 26)

18. The method of claim 14 or 15 wherein the zwitterionic phospholipid is dipalmitoyl phosphatidyl choline.

5 19. The method of claim 14 or 15 wherein the composition further comprises a neutral lipid.

10 20. The method of claim 20 wherein the neutral lipid is a triglyceride.

15 21. A pharmaceutical preparation comprising a non-covalently associated combination of nonsteroidal anti-inflammatory agent, zwitterionic phospholipid, and neutral lipid.

22. The pharmaceutical preparation of claim 24 wherein the zwitterionic phospholipid is dipalmitoyl phosphatidyl choline and the neutral lipid is tripalmitin.

20 23. The pharmaceutical preparation of claim 21 further defined as essentially free of anionic phospholipid.

1/23




Fig. 1A




Fig. 1B

SUBSTITUTE SHEET (RULE 26)

2/23




Fig. 2A




Fig. 2B

SUBSTITUTE SHEET (RULE 26)

3/23




Fig. 3

SUBSTITUTE SHEET (RULE 26)




Fig. 4A




Fig. 4B




Fig. 4C

SUBSTITUTE SHEET (RULE 26)

5/23




Fig. 5

6/23




Fig. 6

SUBSTITUTE SHEET (RULE 26)

7/23




Fig. 7

8/23




Fig. 3

SUBSTITUTE SHEET (RULE 26)

9/23




Fig. 9

SUBSTITUTE SHEET (RULE 26)

10/23




Fig. 10

11/23




Fig. 11

12/23




Fig. 12





Fig. 13



ASA to DPPC Ratios

Fig. 14

= 5/grp





Fig. 15

SUBSTITUTE SHEET (RULE 26)

16/23



SUBSTITUTE SHEET (RULE 26)

17/23





Fig. 17

SUBSTITUTE SHEET (RULE 26)



19/23




Fig. 19

SUBSTITUTE SHEET (RULE 26)




Fig. 20




Fig. 21




Fig. 22




Fig. 23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/00633**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 31/74

US CL : 424/78.05; 514/78, 171

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : A61K 31/74; 424/78.05; 514/78, 171

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
APSElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
NON-STEROIDAL ANTI-INFLAMMATORY DRUGS; PHOSPHOLIPIDS; PHOSPHATIDYL-CHOLINE; SALICYLATE;
ASPIRIN, NAPROXEN, INDOMETHACIN, DICLOFENAC; ANTI-PYRETIC**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LICHTENBERGER et al. "Non-steroidal anti-inflammatory drugs (NSAIDS) associate with zwitterionic phospholipids: Insight into the mechanism and reversal of NSAID-induced gastrointestinal injury", NATURE MEDICINE, Vol. 1, #2, 02/1995. See entire document.	1-23
X	US, A, 4,309,420 (GHYCZY ET AL.), 05 January 1982. See entire document.	1-23
X	US, A, 4,369,182 (GHYCZY ET AL.), 18 January 1983, see entire document.	1-23



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

11 APRIL 1996

Date of mailing of the international search report

27 JUN 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SAMUEL A. ACQUAH aco

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/00633

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4,421,747 (GHYCZY ET AL.) 20 December 1983, see entire document.	1-23
Y	JP, A, 3,176,425 (NIPPON SHINYAKU) 31 July 1991. See the abstract.	1-7, 12, 13, 16